



Question 7: The leaves of an electroscope are initially apart. A positively charged rod is brought near to the top of the electroscope and the leaves of the electroscope fall together.

- A) The electroscope must be negatively charged.
- B) The electroscope isn't charged at all.
- C) The electroscope must be positively charged.
- D) It's impossible to determine if it's positively or negatively charged.

Question 8: The electric field from a point charge is shown at two points along the x-axis. At what position does the charge rest?




Question 9: A water molecule falls from a tap near a charged wand. What happens to the water molecule at the instant shown in the figure?



- A) It rotates counter-clockwise and moves towards the wand.
- B) It rotates clockwise and moves towards the wand.
- C) It rotates counter-clockwise and moves away from the wand.
- D) It rotates clockwise and moves away from the wand.
- E) It doesn't rotate and it moves away from the wand.
- F) It doesn't rotate and it moves towards the wand.


Question 10: The diagram below shows a charged spherical conductor of radius R . The radius of the sphere is doubled to $2R$ and the charge is halved to $Q/2$. What is the ratio of the electric field at point p after the change compared with before the change $E_{\text{after}}/E_{\text{before}}$?

- A) 2
- B) 1
- C) 1/2
- D) 1/4
- E) 1/8




Question 11: The graph shows electric potential along the x-axis. A 20 nC particle is shot from the left ($x < 0$) with 10^{-6} J of kinetic energy. The particle will turn around at


- A) 11 cm
- B) 10 cm
- C) 8 cm
- D) 5 cm
- E) 2 cm
- F) 1 cm
- G) no turning point



Question 12: The diagram to the right shows two thin charged spherical shells. The inner shell has a charge Q and the outer shell has a charge $-Q$.




Which plot below best describes the outward electric field as a function of radius in this situation?



Hedonism-bot will only sit on a platform suspended by pure oxygen. Hedonism-bot has one problem, though. When the temperature outside changes, the gas expands and contracts, which moves the platform up and down and drives Hedonism-bot crazy.

To fix the problem, Hedonism-bot has charged the platform with 1 Coulomb and installed a machine that can generate a uniform electric field of any strength that can point either up or down. When the temperature changes, the electric field will change, thus keeping the platform at a constant height.



If the electric field is initially zero and the temperature drops by 1 degree Kelvin, what will the final value of the electric field need to be?

Possibly useful info: When the temperature is 300 K the platform is 10 m above the bottom of the container. Hedonism-bot has a mass of 1000 kg, and the area of the platform is 5m^2 . The ambient air pressure is 100kPa.

